

LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
5. Total PCB represents the sum of congeners as reported by the laboratory. April 2011 samples were analyzed by Method 1668A; November 2011 samples were analyzed by Method 608.





Cornell-Pohlig Electronics
Superfund Site
South Plainfield, NJ

Total PCB Concentrations in Recently Deposited
Sediments
Boun Brook 04/11/13

2013

FIGURE 6-2a



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Arsenic Concentrations in Recently Deposited
Sediments
Bourne Brook 0-4 F/FS

2013

FIGURE 6-2b





LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.

DRAFT FINAL



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Chromium Concentrations in Recently Deposited
Sediments
Bound Brook 04 R/F/S

2013

FIGURE 6-2d



- LEGEND:**
- ◆ April 2011 Surface Sediment (Be7 bearing)
 - ◇ April 2011 Surface Sediment (non-Be7 bearing)
 - ◆ April 2011 High Resolution Core Top (Be7 bearing)
 - ◆ Nov 2011 Surface Sediment (Be7 bearing)
 - ◆ Nov 2011 Sediment Trap (Be7 bearing)
 - ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

- NOTES:**
1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
 2. For samples with field duplicates, the average concentration is presented.
 3. Nondetected concentrations are presented as half the method detection limit.
 4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Copper Concentrations in Recently Deposited
Sediments
Bourne Brook 0-4 R/F/FS

2013

FIGURE 6-2e



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Lead Concentrations in Recently Deposited Sediments
Bounded Brook 0-4 R/Fs

2013

FIGURE 6-2f



LEGEND:

- April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 High Resolution Core Top (Be7 bearing)
- Nov 2011 Surface Sediment (Be7 bearing)
- Nov 2011 Sediment Trap (Be7 bearing)
- Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

- Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Mercury Concentrations in Recently Deposited
Sediments
Bound Brook 0-4 R/FS

2013

FIGURE 6-2g



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Nickel Concentrations in Recently Deposited Sediments
Boun Brook 0.4 R.F.

2013

FIGURE 6-2h



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Silver Concentrations in Recent Deposited Sediments
Bounded Brook 0.4 R/F

2013

FIGURE 6-2i



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- ◆ Nov 2011 Surface Sediment (Be7 bearing)
- ◆ Nov 2011 Sediment Trap (Be7 bearing)
- ◇ Nov 2011 Sediment Trap (non-Be7 bearing)

NOTES:

1. Filled symbols indicate the presence of Be-7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be-7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Zinc Concentrations in Recently Deposited Sediments
Bounded Brook 0-4 R/Fs

2013

FIGURE 6-2j



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (Be7 bearing, potentially site impacted)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing, potentially site impacted)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Total PAH equals the sum of the 16 priority PAH compounds. Nondetected concentrations were incorporated into the summation as zero.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Total PAH Concentrations in Recently Deposited
Sediments
Bounded Brook 01/4 10/F5

2013

FIGURE 6-2k



LEGEND: <ul style="list-style-type: none">◆ April 2011 Surface Sediment (Be7 bearing)● April 2011 Surface Sediment (Be7 bearing, potentially site impacted)◇ April 2011 Surface Sediment (non-Be7 bearing)○ April 2011 Surface Sediment (non-Be7 bearing, potentially site impacted)◆ April 2011 High Resolution Core Top (Be7 bearing) TOC denotes Total Organic Carbon HMW denotes High Molecular Weight	NOTES: <ol style="list-style-type: none">1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.2. For samples with field duplicates, the average concentration is presented.3. Nondetected concentrations were incorporated into the summation as zero.4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.	





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

High Molecular Weight PAH Concentrations in
Recently Deposited Sediment
Bounded Brook 01/4/13/F5

2013

FIGURE 6-2I



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (Be7 bearing, potentially site impacted)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing, potentially site impacted)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon
LMW denotes Low Molecular Weight

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations were incorporated into the summation as zero.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Low Molecular Weight PAH Concentrations in
Recently Deposited Sediment
Bounded Brook 01/4 10/15

2013

FIGURE 6-2m

DRAFT FINAL





LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.



LEGEND: <ul style="list-style-type: none">◆ April 2011 Surface Sediment (Be7 bearing)◇ April 2011 Surface Sediment (non-Be7 bearing)◆ April 2011 High Resolution Core Top (Be7 bearing)TOC denotes Total Organic Carbon	NOTES: <ol style="list-style-type: none">1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.2. For samples with field duplicates, the average concentration is presented.3. Nondetected concentrations are presented as half the method detection limit.4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.		





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Aldrin Concentrations in Recently Deposited
Sediments
Boun Brook C014 10/15

2013

FIGURE 6-2o



LEGEND:

- April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 High Resolution Core Top (Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

- Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Dieldrin Concentrations in Recently Deposited
Sediments
Boun Brook C014 10/15/13

2013

FIGURE 6-2p



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Endrin Concentrations in Recently Deposited
Sediments
Boun Brook C014 10/F5

2013

FIGURE 6-2q



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

4,4' DDD Concentrations in Recently Deposited
Sediments
Borough of South Plainfield, NJ

2013

FIGURE 6-2r



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

4,4' DDE Concentrations in Recently Deposited
Sediments
Bounded Brook 01/4 10/15

2013

FIGURE 6-2s



LEGEND: <ul style="list-style-type: none">April 2011 Surface Sediment (Be7 bearing)April 2011 Surface Sediment (non-Be7 bearing)April 2011 High Resolution Core Top (Be7 bearing)TOC denotes Total Organic Carbon	NOTES: <ul style="list-style-type: none">1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.2. For samples with field duplicates, the average concentration is presented.3. Nondetected concentrations are presented as half the method detection limit.4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.	





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Alpha-BHC Concentrations in Recently Deposited
Sediments
Boun Brook C014 10/15/13

2013

FIGURE 6-2t



LEGEND:

- April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 High Resolution Core Top (Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

- Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

**Beta-BHC Concentrations in Recently Deposited
Sediments**
Bounded Brook 01/4 10/15

2013

FIGURE 6-2u



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Gamma-BHC Concentrations in Recently Deposited
Sediments
Boun Brook C014 10/15

2013

FIGURE 6-2v





LEGEND:

- April 2011 Surface Sediment (Be7 bearing)
- April 2011 Surface Sediment (non-Be7 bearing)
- April 2011 High Resolution Core Top (Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

- Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Beta-Chlordane Concentrations in Recently
Deposited Sediment
Bounded Brook 01/4 10/15

2013

FIGURE 6-2x



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)

TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.





Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Heptachlor Epoxide Concentrations in Recently
Deposited Sediment
Bounded Brook 04/10/13

2013

FIGURE 6-2y



LEGEND:

- ◆ April 2011 Surface Sediment (Be7 bearing)
- ◇ April 2011 Surface Sediment (non-Be7 bearing)
- ◆ April 2011 High Resolution Core Top (Be7 bearing)
- TOC denotes Total Organic Carbon

NOTES:

1. Filled symbols indicate the presence of Be7 at a concentration greater than 0.5 pCi/g; open symbols indicate a Be7 concentration less than 0.5 pCi/g.
2. For samples with field duplicates, the average concentration is presented.
3. Nondetected concentrations are presented as half the method detection limit.
4. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Hexachlorobenzene Concentrations in Recently
Deposited Sediment
Bounded Brook 01/4 10/F5

2013

FIGURE 6-2z



Photograph Description: Flooding of Bound Brook at Clinton Avenue Bridge before and after storm events – looking north.



Cornell University Center for Environmental and Estuarine Science
Superfund Site
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/FS

2013

FIGURE 6-3a

DRAFT FINAL



Photograph Description: Flooding of Spring Lake outlet before and after storm events - looking upstream from Cedar Brook.



Cornell University Library
Superfund Site
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/F

2013

FIGURE 6-3b

DRAFT FINAL



Photograph Description: Flooding of unnamed tributary near New Brunswick Avenue before and after storm events.



Cornell University Center for Environmental and Estuarine Science
Superfund Site
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/FI

2013

FIGURE 6-3c

DRAFT FINAL



Photograph Description: Flooding of roads from Bound Brook in Cap Lane neighborhood – looking south from north side of brook (brook runs perpendicular to Cap Lane).



Cornell Center for the Environment
Superfund Site
South Plainfield, NJ

**Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee**

Bound Brook OU4 RI/FI

2013

FIGURE 6-3d

DRAFT FINAL



Photograph Description: Flooding of New Market Pond onto roadway near Elsie Road – look north from south side of brook.



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/FI

2013

FIGURE 6-3e

DRAFT FINAL



Photograph Description: Flooding of Bound Brook at New Brunswick Bridge construction site.



Cornell Center for Environmental and Estuarine Science
Superfund Site
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/F

2013

FIGURE 6-3f

DRAFT FINAL



Photograph Description: Debris carried overland into park adjacent to New Market Pond by flooding of Bound Brook and New Market Pond.



Cornell University Center for
Environmental and Estuarine Science
South Plainfield, NJ

Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/F

2013

FIGURE 6-3g

DRAFT FINAL



Photograph Description: New Market Pond Dam at RM3.4 with flow release during Hurricane Irene and Tropical Storm Lee.



Cornell University
Superfund Site
South Plainfield, NJ


Flooding in OU4 Bound Brook Study Area associated
with Hurricane Irene and Tropical Storm Lee

Bound Brook OU4 RI/FI

2013

FIGURE 6-3h

DRAFT FINAL



LEGEND:

- ◆ Contaminant (mg/kg) included in regression
- ◇ Contaminant (mg/kg) not included in regression
- Regression Line

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Regression equation and correlation coefficient calculated by Microsoft Excel.
3. Data sets shown include April 2011 Ekman Samples, November 2011 Ekman Samples, and November 2011 Sediment Trap Samples.




Corbett-Dunbar Electronics
Superfund Site
South Plains River

Iron versus Aluminum Correlation in Recently
Deposited Sediment
Bound Brook OLW R/LFS

2013

FIGURE 6-4a

DRAFT FINAL



LEGEND:

- ◆ Contaminant (mg/kg) included in regression
- Regression Line

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Regression equation and correlaton coefficient calculated by Microsoft Excel.
3. Data sets shown include April 2011 Ekman Samples, November 2011 Ekman Samples, and November 2011 Sediment Trap Samples.



Connecticut Department of Environmental Protection
 State Funded Site
 South Plain Field Station

TOC versus Aluminum Correlation in Recently
 Deposited Sediment
 Bound Brook OLW RIFES

2013


FIGURE 6-4b

DRAFT FINAL

Correlations

	Arsenic	Cadmium	Chromium	Copper	Iron	Lead	Mercury	Nickel	Silver	Zinc
Arsenic	1.00	0.81	0.67	0.60	0.92	0.76	0.68	0.63	0.36	0.85
Cadmium	0.81	1.00	0.77	0.73	0.84	0.83	0.68	0.76	0.51	0.82
Chromium	0.67	0.77	1.00	0.87	0.80	0.86	0.72	0.78	0.56	0.80
Copper	0.60	0.73	0.87	1.00	0.68	0.70	0.51	0.96	0.37	0.65
Iron	0.92	0.84	0.80	0.68	1.00	0.77	0.65	0.69	0.42	0.86
Lead	0.76	0.83	0.86	0.70	0.77	1.00	0.87	0.65	0.72	0.88
Mercury	0.68	0.68	0.72	0.51	0.65	0.87	1.00	0.44	0.75	0.85
Nickel	0.63	0.76	0.78	0.96	0.69	0.65	0.44	1.00	0.27	0.61
Silver	0.36	0.51	0.56	0.37	0.42	0.72	0.75	0.27	1.00	0.68
Zinc	0.85	0.82	0.80	0.65	0.86	0.88	0.85	0.61	0.68	1.00

Scatterplot Matrix




Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Metals Correlation in Recently Deposited
Sediment
Bound Brook OU4 RI/FS

2013

FIGURE 6-5

DRAFT FINAL



LEGEND:

- ◆ Radiological Concentration
- ┃ Uncertainty

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Nondetected concentrations are presented as half the method detection limit.
3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
4. Error bars represent the laboratory reported uncertainty.






Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Radiological Depth Profile in High Resolution
Sediment Core
Bound Brook OL - RI FS

2013




FIGURE 6-6

DRAFT FINAL






LEGEND: ● Contaminant concentration ◆ Radiological concentration TOC denotes Total Organic Carbon	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: ● Contaminant concentration ◆ Radiological concentration	NOTES:
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: <div><div></div> Contaminant concentration</div> <div><div></div> Radiological concentration</div>	NOTES:
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: <div><div>●</div> Contaminant concentration</div> <div><div>◆</div> Radiological concentration</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	








LEGEND: ● Contaminant concentration ◆ Radiological concentration	NOTES:	
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	








LEGEND: <ul style="list-style-type: none">Contaminant concentrationRadiological concentration	NOTES: <ul style="list-style-type: none">1. For samples with field duplicates, the average concentration is presented.2. Nondetected concentrations are presented as half the method detection limit.3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).





LEGEND: ● Contaminant concentration ◆ Radiological concentration	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	








LEGEND:

- Contaminant concentration
- Radiological concentration

NOTES:




- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
- Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: ● Contaminant concentration ◆ Radiological concentration	NOTES:
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).





LEGEND: ● Contaminant concentration ◆ Radiological concentration	NOTES:
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).





LEGEND:

- Contaminant concentration
- Radiological concentration
- TOC denotes Total Organic Carbon

NOTES:

- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
- Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).






Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Contaminant Depth Profile and Geochronology for
Total PAH
Bounded Brook C014 10/F5




2013

FIGURE 6-7k



LEGEND: <ul style="list-style-type: none">Contaminant concentrationRadiological concentrationTOC denotes Total Organic CarbonHMW PAH denotes High Molecular Weight PAHs	NOTES: <ol style="list-style-type: none">For samples with field duplicates, the average concentration is presented.Nondetected concentrations are presented as half the method detection limit.High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).





LEGEND:

- Contaminant concentration
- - -◆- - Radiological concentration
- TOC denotes Total Organic Carbon
- LMW PAH denotes Low Molecular Weight PAHs

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Nondetected concentrations are presented as half the method detection limit.
3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).






Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Contaminant Depth Profile and Geochronology for
LMW PAH
Boulevard Brook C014 10/F5

2013

FIGURE 6-7m



LEGEND:

- Contaminant concentration
- Radiological concentration
- TOC denotes Total Organic Carbon

NOTES:

- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
- Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).






Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Contaminant Depth Profile and Geochronology for
2,3,7,8-TCDD
Bounded Brook C014 10/F5

2013




FIGURE 6-7n

DRAFT FINAL



LEGEND: <div><div><div></div></div><div>Contaminant concentration</div></div> <div><div><div></div></div><div>Radiological concentration</div></div> <div>TOC denotes Total Organic Carbon</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).		





LEGEND:

- Contaminant concentration
- Radiological concentration
- TOC denotes Total Organic Carbon

NOTES:

- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
- Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).






Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Contaminant Depth Profile and Geochronology for
4,4'-DDD
Boulevard Brook C014 10/F5




2013

FIGURE 6-7p






LEGEND: <div><div></div> Contaminant concentration</div> <div><div></div> Radiological concentration</div> <div>TOC denotes Total Organic Carbon</div>	NOTES:
	<div>1. For samples with field duplicates, the average concentration is presented.</div> <div>2. Nondetected concentrations are presented as half the method detection limit.</div> <div>3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.</div> <div>4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).</div>





LEGEND: ● Contaminant concentration ◆ Radiological concentration TOC denotes Total Organic Carbon	NOTES:
	1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND:

- Contaminant concentration
- Radiological concentration
- TOC denotes Total Organic Carbon

NOTES:




- For samples with field duplicates, the average concentration is presented.
- Nondetected concentrations are presented as half the method detection limit.
- High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent.
- Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: <div><div></div> Contaminant concentration</div> <div><div></div> Radiological concentration</div> <div>TOC denotes Total Organic Carbon</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: <div><div></div> Contaminant concentration</div> <div><div></div> Radiological concentration</div> <div>TOC denotes Total Organic Carbon</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).








LEGEND: ● Contaminant concentration ◆ Radiological concentration TOC denotes Total Organic Carbon	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	






LEGEND: <div><div><div></div></div><div>Contaminant concentration</div></div> <div><div><div></div></div><div>Radiological concentration</div></div> <div>TOC denotes Total Organic Carbon</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	





LEGEND: <div><div></div> Contaminant concentration</div> <div><div></div> Radiological concentration</div> <div>TOC denotes Total Organic Carbon</div>	NOTES: 1. For samples with field duplicates, the average concentration is presented. 2. Nondetected concentrations are presented as half the method detection limit. 3. High resolution core top represents the average of the first two slices (0-6 cm total) since both slices were Be7 bearing and had a moisture content of approximately 70 percent. 4. Due to the uncertainty in extrapolating sedimentation rate to the core bottom, depositional horizons are only presented from 2011 to 1925 (or for the top 74 cm of the core).	





LEGEND:

- Natural Log Total PCB (absolute)
- ◆ Natural Log Total PCB (TOC-normalized)
- Linear Regression (Natural Log Total PCB [absolute])
- Linear Regression (Natural Log Total PCB [TOC-normalized])

NOTES:

1. Half-life is calculated by plotting the natural logarithmic Total PCB concentrations (absolute and TOC-normalized) against depositional year and fitting a linear regression. The half-life is equal to 0.693 divided by the slope of the regression.
2. Half-life for Total PCB (absolute) is approximately 40 years.
3. Half-life for Total PCB (TOC-normalized) is approximately 70 years.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ


Estimated Half-Life Calculation for Total PCB
Bound Brook OUI RI/FS

2013


FIGURE 6-8

DRAFT FINAL


**Talmadge Road Bridge
(RM8.3)**




**Belmont Ave Bridge
(RM6.8)**




**Downstream of Site
(RM6.6 to RM0)**




**Green Brook
(Downstream)**



**Green Brook
(Upstream)**



Tributaries



LEGEND:

- Aroclor 1254 Percentage
- Aroclor 1242 Percentage
- Aroclor 1260 Percentage

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Aroclor concentration reported as sum of congeners by either Method 1668 or Method 608.




DRAFT FINAL


2013

FIGURE 6-9


**Talmadge Road Bridge
(RM8.3)**




**Belmont Ave Bridge
(RM6.8)**




**Downstream of Site
(RM6.6 to RM0)**




**Green Brook
(Downstream)**



**Green Brook
(Upstream)**



Tributaries



LEGEND:

- DiCB + TriCB
- TetraCB
- PentaCB
- HeptaCB
- HexaCB
- OctaCB + NonaCB

CB denotes chlorobiphenyl homologue groups

NOTES:

1. For samples with field duplicates, the average concentration is presented.
2. Total PCB concentration is calculated as the sum of the Dichlorobiphenyl through Nonachlorobiphenyl homologues.
3. Nondetected concentrations are not included in the Total PCB summation.
4. Homologue concentrations are reported as sum of congeners by either Method 1668 or Method 608.





Corning Dubner Electronics
Superfund Site
South Plainfield, NJ

Comparison of PCB Homologue Distribution in
Surface Sediments
Bound Brook 04 R/F/S

2013

FIGURE 6-10

DRAFT FINAL



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)
- ED TPCB = Ekman Dredge Total PCB concentration
- ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Surface Sediment Homologue Patterns in
Recently Deposited Sediments


Bound Brook OU4 RI/F

2013


FIGURE 6-11a

DRAFT FINAL

Unnamed Tributary near New Brunswick Avenue, RM 4.66



Downstream of Spring Lake Outfall
(collected on Cedar Brook) RM 6



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)
- ED TPCB = Ekman Dredge Total PCB concentration
- ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ



Surface Sediment Homologue Patterns in
Recently Deposited Sediments

Bourne Brook OU4 RI/F

2013

FIGURE 6-11b

DRAFT FINAL



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)
- ED TPCB = Ekman Dredge Total PCB concentration
- ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ



**Surface Sediment Homologue Patterns in
Recently Deposited Sediments**

Bound Brook OU4 RI/F

2013

FIGURE 6-11c

DRAFT FINAL



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)
- ED TPCB = Ekman Dredge Total PCB concentration
- ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ



Surface Sediment Homologue Patterns in
Recently Deposited Sediments

Bound Brook OU4 RI/F

2013

FIGURE 6-11d

DRAFT FINAL



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)

ED TPCB = Ekman Dredge Total PCB concentration

ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Surface Sediment Homologue Patterns in
Recently Deposited Sediments


Bound Brook OU4 RI/F

2013


FIGURE 6-11e

DRAFT FINAL

Bound Brook RM 4.83



Bound Brook RM4.75



LEGEND:

■ Ekman Dredge (ED)
 TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
 Superfund Site
 South Plainfield, NJ

Surface Sediment Homologue Patterns in
 Recently Deposited Sediments


Bound Brook OU4 RI/F

2013


FIGURE 6-11f

DRAFT FINAL

Bound Brook RM4.38 (April 2011)



Bound Brook RM 4.38 (November 2011)



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)

ED TPCB = Ekman Dredge Total PCB concentration

ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Surface Sediment Homologue Patterns in
Recently Deposited Sediments

Bound Brook OU4 RI/F



2013

FIGURE 6-11g

DRAFT FINAL

Bound Brook RM2.85

Bound Brook RM2.59



LEGEND:

■ Ekman Dredge (ED)
 TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics Corporation
 Superfund Site
 South Plainfield, NJ

Surface Sediment Homologue Patterns in
 Recently Deposited Sediments


Bound Brook OU4 RI/F

2013


FIGURE 6-11h

DRAFT FINAL

Bound Brook RM3.44



Bound Brook RM 2.2



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)

ED TPCB = Ekman Dredge Total PCB concentration

ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Surface Sediment Homologue Patterns in
Recently Deposited Sediments

Bound Brook OU4 RI/F



2013

FIGURE 6-11i

DRAFT FINAL

Bound Brook RM1.29

Bound Brook RM0.15



LEGEND:

■ Ekman Dredge (ED)
 TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
 Superfund Site
 South Plainfield, NJ



Surface Sediment Homologue Patterns in
 Recently Deposited Sediments

Bound Brook OU4 RI/F

2013

FIGURE 6-11j

DRAFT FINAL



LEGEND:

- Ekman Dredge (ED)
- Sediment Trap (ST)

ED TPCB = Ekman Dredge Total PCB concentration

ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell University
Superfund Site
South Plainfield, NJ

Surface Sediment Homologue Patterns in
Recently Deposited Sediments


Bound Brook OU4 RI/FI

2013


FIGURE 6-11k

DRAFT FINAL

Green Brook - Downstream of Confluence
with Bound Brook, RM -0.1



Green Brook RM-1.5



LEGEND:

■ Ekman Dredge (ED)
■ Sediment Trap (ST)
ED TPCB = Ekman Dredge Total PCB concentration
ST TPCB = Sediment Trap Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. Total PCB concentration was rounded to 2 or 3 significant digits.
4. Data source: 2011 OU4 Remedial Investigation



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ



Surface Sediment Homologue Patterns in
Recently Deposited Sediments

Bound Brook OU4 RI/F

2013

FIGURE 6-11I

DRAFT FINAL



LEGEND:

Homologue loading factor

NOTES:

1. Only Be7-bearing sediments collected via Ekman dredge or sediment trap are presented, except one Woodbrook sample at RM7.85.
2. Mono- to Deca-chlorobiphenyl homologues were included in the analysis. Homologues reported by Method 1668A or Method 608.
3. Nondetected concentrations were incorporated as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Sources: 2011 OU4 Remedial Investigation and 2009 Woodbrook Road Dump Superfund Site Addendum to Draft Site Characterization Report.




Cornell-Dubilier
Electronics Superfund Site
South Plainfield, NJ

**Principal Components Loading for PCB
Homologues in Recently Deposited Sediment**
Boulevard, OU4, RI/FS

2013

FIGURE 6-12

DRAFT FINAL



LEGEND:

- Bound Brook (April 2011)
 - Bound Brook and Green Brook (November 2011)
 - Tributaries and Upstream Locations (November 2011 and 2009)
 - Downstream of Spring Lake (November 2011)
- UGB = Green Brook upstream of confluence with Bound Brook
 NB = Unnamed tributary near New Brunswick Avenue
 EL = Unnamed tributary near Elsie Avenue
 SL = Downstream of Spring Lake outfall (collected on Cedar Brook)

NOTES:

1. Numbers indicate Bound Brook river mile.
2. Letters indicate geographical location.
3. Only Be7-bearing sediments collected via Ekman dredge or sediment trap are presented, except one Woodbrook sample at RM7.85.
4. Mono- to Deca-chlorobiphenyl homologues were included in the analysis. Homologues reported by Method 1668A or Method 608.
5. Nondetected concentrations were incorporated as zero.
6. Field duplicates are presented as an average concentration.
7. Data Sources: 2011 OU4 Remedial Investigation and 2009 Woodbrook Road Dump Superfund Site Addendum to Draft Site Characterization Report.



Cornell-Dubilier
 Electronics Superfund Site
 South Plainfield, NJ


Principal Component Analysis Results for PCB Homologues in Recently Deposited Sediment

Bound Brook, NJ

2013

FIGURE 6-13

DRAFT FINAL



LEGEND:

- ◇ Bound Brook, Fish Carcass Tissue
- ◆ Bound Brook, Fish Fillet Tissue
- △ Spring Lake, Fish Carcass Tissue
- ▲ Spring Lake, Fish Fillet Tissue

NOTES:

1. Tissue data represent detected contaminant concentrations only.
2. Data represent several fish species (collected in 1997), including blue gill sunfish, carp, catfish, largemouth bass, pumpkinseed sunfish, and white sucker.
3. Spring Lake data projected at river mile 6.2 (note that all fish were caught in Spring Lake, not on Bound Brook).
4. Data source: 1999 USEPA OU4 Ecological Evaluation Report



DRAFT FINAL


Corning Dubner Electronics
Superfund Site
South Plainfield, NJ

Total PCB Concentrations Detected in Fish Tissue
in 1997
Bound Brook OU4 R/F/S


2013

FIGURE 6-14

RM 7.86; Pumpkinseed



RM 7.86; White Sucker



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ RI/F

2013


FIGURE 6-15a

DRAFT FINAL

RM 6.5; Pumpkinseed



RM 6.5; White Sucker



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

- Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
- Nondetected concentrations were incorporated as zero.
- For samples with field duplicates, the average concentration is presented.
- Total PCB concentration was rounded to 2 significant digits.
- Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ

2013


FIGURE 6-15b

DRAFT FINAL

RM 6.5; White Sucker



RM 6.32; Pumpkinseed



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ



Fish Fillet Homologue Patterns

Bound Brook, NJ RI/F

2013

FIGURE 6-15c

DRAFT FINAL



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ RI/F

2013


FIGURE 6-15d

DRAFT FINAL

RM 5.66; White Sucker



RM 5.19; Pumpkinseed



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

- Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
- Nondetected concentrations were incorporated as zero.
- For samples with field duplicates, the average concentration is presented.
- Total PCB concentration was rounded to 2 significant digits.
- Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ RI/F

2013


FIGURE 6-15e

DRAFT FINAL

RM 5.19; Blue Gill Sunfish



RM 5.19; White Sucker



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ/RI/PA

2013


FIGURE 6-15f

DRAFT FINAL

RM 3.71; White Sucker



RM 3.71; Carp



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook, NJ

2013

FIGURE 6-15g

DRAFT FINAL

RM 3.71; Blue Gill Sunfish



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

- Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
- Nondetected concentrations were incorporated as zero.
- For samples with field duplicates, the average concentration is presented.
- Total PCB concentration was rounded to 2 significant digits.
- Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ

Fish Fillet Homologue Patterns


Bound Brook DU/RI/F

2013


FIGURE 6-15h

DRAFT FINAL



Spring Lake; Blue Gill Sunfish



Spring Lake; White Sucker



LEGEND:

 Homologue mass fraction in fish fillet sample
 Homologue mass fraction in fish carcass sample
 TPCB = Total PCB Concentration
 Sample label on top of graphic: "RM 7.86; Pumpkinseed"
 Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ


Fish Fillet Homologue Patterns

Bound Brook, NJ RI/F

2013

FIGURE 6-15i

DRAFT FINAL



LEGEND:

- Homologue mass fraction in fish fillet sample
- Homologue mass fraction in fish carcass sample

TPCB = Total PCB Concentration

Sample label on top of graphic: "RM 7.86; Pumpkinseed"

Indicates River Mile in Bound Brook and Fish Species

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 significant digits.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier Electronics
Superfund Site
South Plainfield, NJ



Fish Fillet Homologue Patterns

Bound Brook DU/RI/F

2013

FIGURE 6-15j

DRAFT FINAL



LEGEND:

■ PCB loading factor

NOTES:

1. Only fish fillet was included in the analysis.
2. 118 congeners were included in the analysis.
3. Nondetected concentrations were incorporated as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.




Cornell-Dubilier
Electronics Superfund Site
South Plainfield, NJ

**Principal Components Loading for PCB
Congeners in Fish Fillets**
Bound Brook, NJ

2013

FIGURE 6-16

DRAFT FINAL



LEGEND:

- New Market Pond
- RM 4.09 to RM 5.22
- RM 5.22 to RM 6.18
- RM 6.18 to RM 6.82
- RM 6.82 to RM 8.31
- Spring Lake

NOTES:

1. Letters indicate fish species:
BG = Blue Gill Sunfish
C = Carp
P = Pumpkinseed Sunfish
WS = White Sucker
2. Only fish fillet was included in the analysis.
3. 118 congeners were included in the analysis.
4. Nondetected concentrations were incorporated as zero.
5. For samples with field duplicates, the average concentration is presented.
6. Data source: 2010 Cornell-Dubilier Bound Brook Reassessment.



Cornell-Dubilier
Electronics Superfund Site
South Plainfield, NJ


Principal Components Analysis Results for PCB
Congeners in Fish Fillets
Bound Brook, NJ

2013


FIGURE 6-17

DRAFT FINAL

RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L



RM6.56; ERT-1; 07/13/10; 46-56 ft; TPCB = 0.0033 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site

Bourne Brook OU3 RI/FI



2013

FIGURE 6-18a

DRAFT FINAL

RM6.55; MW-05; 04/08/10; 25-45.5 ft; TPCB = 14 ug/L

RM6.55; MW-05; 07/14/10; 25-45.5 ft; TPCB = 5.4 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013

FIGURE 6-18b

DRAFT FINAL

RM6.47; MW-11; 07/14/10; 34-59 ft; TPCB = 320 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell University
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18c

DRAFT FINAL

RM6.47; MW-14D; 04/06/10; 199-209 ft; TPCB = 3.5 ug/L



RM6.47; MW-14D; 07/13/10; 199-209 ft; TPCB = 2.3 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18d

DRAFT FINAL

RM6.47; MW-14S; 04/06/10; 65-70 ft; TPCB = 68,000 ug/L



RM6.47; MW-14S; 07/13/10; 65-70 ft; TPCB = 81,000 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/F

2013


FIGURE 6-18e

DRAFT FINAL

RM6.47; MW-15D; 04/06/10; 125-135 ft;
TPCB = 0.36 ug/L



RM6.47; MW-15D; 07/13/10; 125-135 ft;
TPCB = 0.11 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18f

DRAFT FINAL

RM6.47; MW-08; 03/25/10; 42-57.5 ft; TPCB = 10 ug/L



RM6.47; MW-08; 07/14/10; 42-57.5 ft; TPCB = 12 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/F

2013

FIGURE 6-18g

DRAFT FINAL

RM6.47; MW-11; 07/14/10; 205-215 ft; TPCB = 320 ug/L -



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell University
Department of Environmental Engineering
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/F

2013


FIGURE 6-18h

DRAFT FINAL

RM6.44; MW-03; 03/25/10; 17-32 ft; TPCB = 10 ug/L



RM6.44; MW-03; 07/12/10; 17-32 ft; TPCB = 7 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18i

DRAFT FINAL

RM6.44; MW-12; 04/05/10; 35-60 ft;
TPCB = 1,500 ug/L



RM6.44; MW-12; 07/13/10; 35-60 ft;
TPCB = 220 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located on the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18j

DRAFT FINAL

RM6.42; FPW; 04/07/10; 31-41 ft; TPCB = 1.4 ug/L



RM6.42; FPW; 07/12/10; 31-41 ft; TPCB = 0.16 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/F

2013


FIGURE 6-18k

DRAFT FINAL


RM6.42; FPW; 04/07/10; 300-310 ft; TPCB = 0.2 ug/L



RM6.42; FPW; 07/12/10; 300-310 ft; TPCB = 0.01 ug/L



LEGEND:

 Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located on the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18I

DRAFT FINAL

RM6.18; MW-01A; 03/24/10; 24-49 ft; TPCB = 1.1 ug/L



RM6.18; MW-01A; 07/12/10; 24-49 ft; TPCB = 1.3 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/F

2013


FIGURE 6-18m

DRAFT FINAL

RM6.18; MW-16; 04/12/10; 195-205 ft; TPCB = 0.15 ug/L



RM6.18; MW-16; 07/12/10; 195-205 ft; TPCB = 0.42 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Well Located in the Former CDE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18n

DRAFT FINAL

RM6.16; ERT-2; 03/31/10; 25-35 ft;
TPCB = 0.37 ug/L



RM6.16; ERT-2; 07/14/10; 25-35 ft;
TPCB = 0.82 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell University
Superfund Site
South Plainfield, NJ


Groundwater PCB Homologue Patterns for
Wells Outside the Former DOE Site

Bourne Brook OU3 RI/F


2013

FIGURE 6-18o

DRAFT FINAL



RM6.16; MW-17; 03/26/10; 205-215 ft;
 TPCB = 0.09 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
 Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
 Superfund Site
 South Plainfield, NJ


Groundwater PCB Homologue Patterns for
 Wells Outside the Former DOE Site

Bourne Brook OU3 RI/FI


2013

FIGURE 6-18q

DRAFT FINAL



RM5.99; ERT-6; 03/26/10; 75-85 ft; TPCB = 0.0036 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Outside the Former DOE Site


Bourne Brook OU3 RI/F

2013


FIGURE 6-18s

DRAFT FINAL

RM6; ERT-4; 03/25/10; 83-88 ft; TPCB = 0.01 ug/L



RM6; ERT-4; 07/12/10; 83-88 ft; TPCB = 0.05 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Outside the Former DOE Site

Bourne Brook OU3 RI/F


2013

FIGURE 6-18t

DRAFT FINAL


RM6.39; MW-13; 04/07/10; 230-240 ft;

TPCB = 0.12 ug/L



RM6.39; MW-13; 07/14/10; 230-240 ft;

TPCB = 01 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Outside the Former DOE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18u

DRAFT FINAL


RM6.56; ERT-8; 04/01/10; 87-97 ft;
TPCB = 0.0038 ug/L



RM6.56; ERT-8; 07/14/10; 87-97 ft;
TPCB = 0.021 ug/L



LEGEND:

 Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Outside the Former DOE Site


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18v

DRAFT FINAL

RM6.02; MW-20; 03/30/10; 85-95 ft;
TPCB = 0.03 ug/L



RM6.02; MW-20; 07/12/10; 85-95 ft;
TPCB = 0.14 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Located in Spring Lake Area


Bourne Brook OU3 RI/FI

2013


FIGURE 6-18w

DRAFT FINAL

RM5.9; MW-19; 03/29/10; 132-142 ft;
TPCB = 0.02 ug/L



RM5.9; MW-19; 0713/10; 132-142 ft;
TPCB = 0.35 ug/L



LEGEND:

Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell University
Department of Environmental Engineering
Superfund Site
South Plainfield, NJ

Groundwater PCB Homologue Patterns for
Wells Located in Spring Lake Area


Bourne Brook OU3 RI/F

2013


FIGURE 6-18x

DRAFT FINAL


RM6.26; MW-22; 03/5/10; 45-55 ft;
TPCB = 0.000066 ug/L



RM6.26; MW-22; 07/12/10; 45-55 ft;
TPCB = 0.028 ug/L



LEGEND:

 Homologue mass fraction in groundwater sample

Sample Label on top of graphic "RM6.56; ERT-1; 04/08/10; 46-56 ft; TPCB = 0.0033 ug/L"
Indicates River Mile; Well Name; Sample collection date; depth of screen; Total PCB concentration.

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Total PCB concentration was rounded to 2 or 3 significant digits.
3. Nondetected concentrations were included as zero.
4. For samples with field duplicates, the average concentration is presented.
5. Data Source: 2010 OU3 Groundwater RI data



Cornell Dubilier Electronics Corporation
Superfund Site
South Plainfield, NJ


Groundwater PCB Homologue Patterns for
Off-site Wells

Bourne Brook OU3 RI/F

2013

FIGURE 6-18y

DRAFT FINAL




Cornell-Dubilier
Electronics Superfund Site
South Plainfield, NJ

Sediment, Fish, and Groundwater PCB Molecular
Weight Across the OU4 Bound Brook Study Area
Bound Brook RI/FS


2013
FIGURE 6-19

DRAFT FINAL


Depth = 0-6 cm
Approximate Year = 2011



Depth = 6-11 cm
Approximate Year = 2005



LEGEND:

 Homologue fraction in high resolution core slice
TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 or 3 significant digits.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell Dubilier Electronics
Superfund Site
South Plainfield, NJ

PCB Homologue Patterns in High Resolution



Sediment Core

Bourne Brook OU4 RI/F

2013

FIGURE 6-20a

DRAFT FINAL



LEGEND:

- Homologue fraction in high resolution core slice
- TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 or 3 significant digits.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell University
Superfund Site
South Plainfield, NJ

PCB Homologue Patterns in High Resolution



Sediment Core

Bourne Brook OU4 RI/F


2013

FIGURE 6-20b

DRAFT FINAL



LEGEND:

 Homologue fraction in high resolution core slice

TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 or 3 significant digits.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell University
Department of Civil and Environmental Engineering
Superfund Site
South Plainfield, NJ

PCB Homologue Patterns in High Resolution

Sediment Core


Bourne Brook OU4 RI/F

2013


FIGURE 6-20c

DRAFT FINAL

Depth = 31-36 cm
Approximate Year = 1959



Depth = 36-41 cm
Approximate Year = 1956



LEGEND:

■ Homologue fraction in high resolution core slice
TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 or 3 significant digits.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell University
Superfund Site
South Plainfield, NJ

PCB Homologue Patterns in High Resolution

Sediment Core


Bourne Brook OU4 RI/F

2013


FIGURE 6-20d

DRAFT FINAL

Depth = 41-46 cm
Approximate Year = 1954



LEGEND:

 Homologue fraction in high resolution core slice
TPCB = Total PCB concentration

NOTES:

1. Homologue mass fraction (ratio of homologue to Total PCB) was plotted.
2. Nondetected concentrations were incorporated as zero.
3. For samples with field duplicates, the average concentration is presented.
4. Total PCB concentration was rounded to 2 or 3 significant digits.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell University
Superfund Site
South Plainfield, NJ

PCB Homologue Patterns in High Resolution


Sediment Core

Bourne Brook OU4 RI/F

2013

FIGURE 6-20e

DRAFT FINAL



LEGEND:



High resolution sediment core sample

NOTES:

1. For samples with field duplicates, the average concentration is presented..
2. Nondetected concentrations were incorporated as zero.
3. Data represents the PCB molecular weight calculated for surface and subsurface samples from the high resolution sediment core. Only samples above 46 cm or the 1954 depositional horizon are presented.
4. The top slice (0-6 cm) represents the average of two slices since they were both Be7 bearing and represent 2011.
5. Data source: 2011 OU4 Remedial Investigation.



Cornell-Dubilier
Electronics Superfund Site
South Plainfield, NJ

PCB Molecular Weight for Depositional
Year in High Resolution Sediment Core
Bou OU4 RI/FS

2013

FIGURE 6-21

DRAFT FINAL